

Water quality — Guidance on the scope and selection of fish sampling methods

The European Standard EN 14962:2006 has the status of a
British Standard

ICS 13.060.70

National foreword

This British Standard is the official English language version of EN 14962:2006.

The UK participation in its preparation was entrusted by Technical Committee EH/3, Water quality, to Subcommittee EH/3/5, Biological methods, which has the responsibility to:

- aid enquirers to understand the text;
- present to the responsible international/European committee any enquiries on the interpretation, or proposals for change, and keep UK interests informed;
- monitor related international and European developments and promulgate them in the UK.

A list of organizations represented on this subcommittee can be obtained on request to its secretary.

Cross-references

The British Standards which implement international or European publications referred to in this document may be found in the *BSI Catalogue* under the section entitled “International Standards Correspondence Index”, or by using the “Search” facility of the *BSI Electronic Catalogue* or of British Standards Online.

This publication does not purport to include all the necessary provisions of a contract. Users are responsible for its correct application.

Compliance with a British Standard does not of itself confer immunity from legal obligations.

Summary of pages

This document comprises a front cover, an inside front cover, the EN title page, pages 2 to 25 and a back cover.

The BSI copyright notice displayed in this document indicates when the document was last issued.

Amendments issued since publication

Amd. No.	Date	Comments

This British Standard was published under the authority of the Standards Policy and Strategy Committee on 31 May 2006

© BSI 2006

ISBN 0 580 48346 0

National foreword

This British Standard is the official English language version of EN 14962:2006.

The UK participation in its preparation was entrusted by Technical Committee EH/3, Water quality, to Subcommittee EH/3/5, Biological methods, which has the responsibility to:

- aid enquirers to understand the text;
- present to the responsible international/European committee any enquiries on the interpretation, or proposals for change, and keep UK interests informed;
- monitor related international and European developments and promulgate them in the UK.

A list of organizations represented on this subcommittee can be obtained on request to its secretary.

Cross-references

The British Standards which implement international or European publications referred to in this document may be found in the *BSI Catalogue* under the section entitled “International Standards Correspondence Index”, or by using the “Search” facility of the *BSI Electronic Catalogue* or of British Standards Online.

This publication does not purport to include all the necessary provisions of a contract. Users are responsible for its correct application.

Compliance with a British Standard does not of itself confer immunity from legal obligations.

Summary of pages

This document comprises a front cover, an inside front cover, the EN title page, pages 2 to 25 and a back cover.

The BSI copyright notice displayed in this document indicates when the document was last issued.

Amendments issued since publication

Amd. No.	Date	Comments

This British Standard was published under the authority of the Standards Policy and Strategy Committee on 31 May 2006

© BSI 2006

ISBN 0 580 48346 0

National foreword

This British Standard is the official English language version of EN 14962:2006.

The UK participation in its preparation was entrusted by Technical Committee EH/3, Water quality, to Subcommittee EH/3/5, Biological methods, which has the responsibility to:

- aid enquirers to understand the text;
- present to the responsible international/European committee any enquiries on the interpretation, or proposals for change, and keep UK interests informed;
- monitor related international and European developments and promulgate them in the UK.

A list of organizations represented on this subcommittee can be obtained on request to its secretary.

Cross-references

The British Standards which implement international or European publications referred to in this document may be found in the *BSI Catalogue* under the section entitled “International Standards Correspondence Index”, or by using the “Search” facility of the *BSI Electronic Catalogue* or of British Standards Online.

This publication does not purport to include all the necessary provisions of a contract. Users are responsible for its correct application.

Compliance with a British Standard does not of itself confer immunity from legal obligations.

Summary of pages

This document comprises a front cover, an inside front cover, the EN title page, pages 2 to 25 and a back cover.

The BSI copyright notice displayed in this document indicates when the document was last issued.

Amendments issued since publication

Amd. No.	Date	Comments

This British Standard was published under the authority of the Standards Policy and Strategy Committee on 31 May 2006

© BSI 2006

ISBN 0 580 48346 0

Foreword

This document (EN 14962:2006) has been prepared by Technical Committee CEN/TC 230 "Water analysis", the secretariat of which is held by DIN.

This European Standard shall be given the status of a national standard, either by publication of an identical text or by endorsement, at the latest by October 2006, and conflicting national standards shall be withdrawn at the latest by October 2006.

According to the CEN/CENELEC Internal Regulations, the national standards organizations of the following countries are bound to implement this European Standard: Austria, Belgium, Cyprus, Czech Republic, Denmark, Estonia, Finland, France, Germany, Greece, Hungary, Iceland, Ireland, Italy, Latvia, Lithuania, Luxembourg, Malta, Netherlands, Norway, Poland, Portugal, Romania, Slovakia, Slovenia, Spain, Sweden, Switzerland and United Kingdom.

Introduction

This document is developed to select appropriate fish sampling methods for the evaluation of the species composition, abundance and age structure of fish in rivers, lakes and transitional waters.

WARNING — Persons using this document should be familiar with normal laboratory practice. This document does not purport to address all of the safety problems, if any, associated with its use. It is the responsibility of the user to establish appropriate safety and health practices and to ensure compliance with any national regulatory conditions.

IMPORTANT — It is absolutely essential that tests conducted according to this document be carried out by suitably trained staff.

Introduction

This document is developed to select appropriate fish sampling methods for the evaluation of the species composition, abundance and age structure of fish in rivers, lakes and transitional waters.

WARNING — Persons using this document should be familiar with normal laboratory practice. This document does not purport to address all of the safety problems, if any, associated with its use. It is the responsibility of the user to establish appropriate safety and health practices and to ensure compliance with any national regulatory conditions.

IMPORTANT — It is absolutely essential that tests conducted according to this document be carried out by suitably trained staff.

Introduction

This document is developed to select appropriate fish sampling methods for the evaluation of the species composition, abundance and age structure of fish in rivers, lakes and transitional waters.

WARNING — Persons using this document should be familiar with normal laboratory practice. This document does not purport to address all of the safety problems, if any, associated with its use. It is the responsibility of the user to establish appropriate safety and health practices and to ensure compliance with any national regulatory conditions.

IMPORTANT — It is absolutely essential that tests conducted according to this document be carried out by suitably trained staff.

Table 2 — Categories for lakes

Category	Presence or absence of a pelagic or profundal zone	Area km ²
Lake category 1	With or without a pelagic or profundal zone	< 0,5
Lake category 2	Without a pelagic or profundal zone	> 0,5
Lake category 3	With a pelagic and profundal zone	> 0,5

Table 3 — Categories for transitional waters

Category	Width m	max. Depth m
Transitional water category 1	< 5	< 1
Transitional water category 2	> 5	< 2
Transitional water category 3	< 30	> 2
Transitional water category 4	30 to 100	> 2
Transitional water category 5	> 100	> 2

Within a river or transitional water, categories according to the Tables 1 and 3 usually change from lower to higher ones according to longitudinal changes. But also within a relatively short stretch of a river or transitional water, categories can change from lower to higher ones and vice versa. In this case, the area itself of a river or transitional water, and not the whole river or transitional water, should be classified according to the given categories.

6 Fish sampling

6.1 General

Sampling methods are separated into methods by capture (6.2) and data sampling without fish capture (6.3). The principles are explained and the target water types or categories and fish species are given. The suitability of the methods is evaluated depending on the different water categories and their habitats as well as the target species. Target species are those species which can efficiently be caught by a particular method. Tables 4 to 6 provide a synopsis on sampling methods suitable or useful for the determination of the population parameters.

Methods not described below may be used in order to obtain additional information. They shall not be used exclusively.

6.2 Fish sampling by capture

6.2.1 Principle of fish capture

Fish may be caught by active gear (6.2.2 to 6.2.4) or passive (6.2.5 to 6.2.10). For all gear seasonal aspects shall be taken into account. The efficiency of active gear (electric fishing, seining, trawling) especially may vary considerably depending on the time (day/night) of use. Active gear is characterized by a comparatively short-term use, whereas passive ones normally are exposed over longer time periods. The fishing devices used shall be adapted to the local conditions as well as to the expected fish community. If it is intended to

Table 2 — Categories for lakes

Category	Presence or absence of a pelagic or profundal zone	Area km ²
Lake category 1	With or without a pelagic or profundal zone	< 0,5
Lake category 2	Without a pelagic or profundal zone	> 0,5
Lake category 3	With a pelagic and profundal zone	> 0,5

Table 3 — Categories for transitional waters

Category	Width m	max. Depth m
Transitional water category 1	< 5	< 1
Transitional water category 2	> 5	< 2
Transitional water category 3	< 30	> 2
Transitional water category 4	30 to 100	> 2
Transitional water category 5	> 100	> 2

Within a river or transitional water, categories according to the Tables 1 and 3 usually change from lower to higher ones according to longitudinal changes. But also within a relatively short stretch of a river or transitional water, categories can change from lower to higher ones and vice versa. In this case, the area itself of a river or transitional water, and not the whole river or transitional water, should be classified according to the given categories.

6 Fish sampling

6.1 General

Sampling methods are separated into methods by capture (6.2) and data sampling without fish capture (6.3). The principles are explained and the target water types or categories and fish species are given. The suitability of the methods is evaluated depending on the different water categories and their habitats as well as the target species. Target species are those species which can efficiently be caught by a particular method. Tables 4 to 6 provide a synopsis on sampling methods suitable or useful for the determination of the population parameters.

Methods not described below may be used in order to obtain additional information. They shall not be used exclusively.

6.2 Fish sampling by capture

6.2.1 Principle of fish capture

Fish may be caught by active gear (6.2.2 to 6.2.4) or passive (6.2.5 to 6.2.10). For all gear seasonal aspects shall be taken into account. The efficiency of active gear (electric fishing, seining, trawling) especially may vary considerably depending on the time (day/night) of use. Active gear is characterized by a comparatively short-term use, whereas passive ones normally are exposed over longer time periods. The fishing devices used shall be adapted to the local conditions as well as to the expected fish community. If it is intended to

Table 2 — Categories for lakes

Category	Presence or absence of a pelagic or profundal zone	Area km ²
Lake category 1	With or without a pelagic or profundal zone	< 0,5
Lake category 2	Without a pelagic or profundal zone	> 0,5
Lake category 3	With a pelagic and profundal zone	> 0,5

Table 3 — Categories for transitional waters

Category	Width m	max. Depth m
Transitional water category 1	< 5	< 1
Transitional water category 2	> 5	< 2
Transitional water category 3	< 30	> 2
Transitional water category 4	30 to 100	> 2
Transitional water category 5	> 100	> 2

Within a river or transitional water, categories according to the Tables 1 and 3 usually change from lower to higher ones according to longitudinal changes. But also within a relatively short stretch of a river or transitional water, categories can change from lower to higher ones and vice versa. In this case, the area itself of a river or transitional water, and not the whole river or transitional water, should be classified according to the given categories.

6 Fish sampling

6.1 General

Sampling methods are separated into methods by capture (6.2) and data sampling without fish capture (6.3). The principles are explained and the target water types or categories and fish species are given. The suitability of the methods is evaluated depending on the different water categories and their habitats as well as the target species. Target species are those species which can efficiently be caught by a particular method. Tables 4 to 6 provide a synopsis on sampling methods suitable or useful for the determination of the population parameters.

Methods not described below may be used in order to obtain additional information. They shall not be used exclusively.

6.2 Fish sampling by capture

6.2.1 Principle of fish capture

Fish may be caught by active gear (6.2.2 to 6.2.4) or passive (6.2.5 to 6.2.10). For all gear seasonal aspects shall be taken into account. The efficiency of active gear (electric fishing, seining, trawling) especially may vary considerably depending on the time (day/night) of use. Active gear is characterized by a comparatively short-term use, whereas passive ones normally are exposed over longer time periods. The fishing devices used shall be adapted to the local conditions as well as to the expected fish community. If it is intended to

Table 2 — Categories for lakes

Category	Presence or absence of a pelagic or profundal zone	Area km ²
Lake category 1	With or without a pelagic or profundal zone	< 0,5
Lake category 2	Without a pelagic or profundal zone	> 0,5
Lake category 3	With a pelagic and profundal zone	> 0,5

Table 3 — Categories for transitional waters

Category	Width m	max. Depth m
Transitional water category 1	< 5	< 1
Transitional water category 2	> 5	< 2
Transitional water category 3	< 30	> 2
Transitional water category 4	30 to 100	> 2
Transitional water category 5	> 100	> 2

Within a river or transitional water, categories according to the Tables 1 and 3 usually change from lower to higher ones according to longitudinal changes. But also within a relatively short stretch of a river or transitional water, categories can change from lower to higher ones and vice versa. In this case, the area itself of a river or transitional water, and not the whole river or transitional water, should be classified according to the given categories.

6 Fish sampling

6.1 General

Sampling methods are separated into methods by capture (6.2) and data sampling without fish capture (6.3). The principles are explained and the target water types or categories and fish species are given. The suitability of the methods is evaluated depending on the different water categories and their habitats as well as the target species. Target species are those species which can efficiently be caught by a particular method. Tables 4 to 6 provide a synopsis on sampling methods suitable or useful for the determination of the population parameters.

Methods not described below may be used in order to obtain additional information. They shall not be used exclusively.

6.2 Fish sampling by capture

6.2.1 Principle of fish capture

Fish may be caught by active gear (6.2.2 to 6.2.4) or passive (6.2.5 to 6.2.10). For all gear seasonal aspects shall be taken into account. The efficiency of active gear (electric fishing, seining, trawling) especially may vary considerably depending on the time (day/night) of use. Active gear is characterized by a comparatively short-term use, whereas passive ones normally are exposed over longer time periods. The fishing devices used shall be adapted to the local conditions as well as to the expected fish community. If it is intended to

Table 2 — Categories for lakes

Category	Presence or absence of a pelagic or profundal zone	Area km ²
Lake category 1	With or without a pelagic or profundal zone	< 0,5
Lake category 2	Without a pelagic or profundal zone	> 0,5
Lake category 3	With a pelagic and profundal zone	> 0,5

Table 3 — Categories for transitional waters

Category	Width m	max. Depth m
Transitional water category 1	< 5	< 1
Transitional water category 2	> 5	< 2
Transitional water category 3	< 30	> 2
Transitional water category 4	30 to 100	> 2
Transitional water category 5	> 100	> 2

Within a river or transitional water, categories according to the Tables 1 and 3 usually change from lower to higher ones according to longitudinal changes. But also within a relatively short stretch of a river or transitional water, categories can change from lower to higher ones and vice versa. In this case, the area itself of a river or transitional water, and not the whole river or transitional water, should be classified according to the given categories.

6 Fish sampling

6.1 General

Sampling methods are separated into methods by capture (6.2) and data sampling without fish capture (6.3). The principles are explained and the target water types or categories and fish species are given. The suitability of the methods is evaluated depending on the different water categories and their habitats as well as the target species. Target species are those species which can efficiently be caught by a particular method. Tables 4 to 6 provide a synopsis on sampling methods suitable or useful for the determination of the population parameters.

Methods not described below may be used in order to obtain additional information. They shall not be used exclusively.

6.2 Fish sampling by capture

6.2.1 Principle of fish capture

Fish may be caught by active gear (6.2.2 to 6.2.4) or passive (6.2.5 to 6.2.10). For all gear seasonal aspects shall be taken into account. The efficiency of active gear (electric fishing, seining, trawling) especially may vary considerably depending on the time (day/night) of use. Active gear is characterized by a comparatively short-term use, whereas passive ones normally are exposed over longer time periods. The fishing devices used shall be adapted to the local conditions as well as to the expected fish community. If it is intended to

Table 2 — Categories for lakes

Category	Presence or absence of a pelagic or profundal zone	Area km ²
Lake category 1	With or without a pelagic or profundal zone	< 0,5
Lake category 2	Without a pelagic or profundal zone	> 0,5
Lake category 3	With a pelagic and profundal zone	> 0,5

Table 3 — Categories for transitional waters

Category	Width m	max. Depth m
Transitional water category 1	< 5	< 1
Transitional water category 2	> 5	< 2
Transitional water category 3	< 30	> 2
Transitional water category 4	30 to 100	> 2
Transitional water category 5	> 100	> 2

Within a river or transitional water, categories according to the Tables 1 and 3 usually change from lower to higher ones according to longitudinal changes. But also within a relatively short stretch of a river or transitional water, categories can change from lower to higher ones and vice versa. In this case, the area itself of a river or transitional water, and not the whole river or transitional water, should be classified according to the given categories.

6 Fish sampling

6.1 General

Sampling methods are separated into methods by capture (6.2) and data sampling without fish capture (6.3). The principles are explained and the target water types or categories and fish species are given. The suitability of the methods is evaluated depending on the different water categories and their habitats as well as the target species. Target species are those species which can efficiently be caught by a particular method. Tables 4 to 6 provide a synopsis on sampling methods suitable or useful for the determination of the population parameters.

Methods not described below may be used in order to obtain additional information. They shall not be used exclusively.

6.2 Fish sampling by capture

6.2.1 Principle of fish capture

Fish may be caught by active gear (6.2.2 to 6.2.4) or passive (6.2.5 to 6.2.10). For all gear seasonal aspects shall be taken into account. The efficiency of active gear (electric fishing, seining, trawling) especially may vary considerably depending on the time (day/night) of use. Active gear is characterized by a comparatively short-term use, whereas passive ones normally are exposed over longer time periods. The fishing devices used shall be adapted to the local conditions as well as to the expected fish community. If it is intended to

Table 2 — Categories for lakes

Category	Presence or absence of a pelagic or profundal zone	Area km ²
Lake category 1	With or without a pelagic or profundal zone	< 0,5
Lake category 2	Without a pelagic or profundal zone	> 0,5
Lake category 3	With a pelagic and profundal zone	> 0,5

Table 3 — Categories for transitional waters

Category	Width m	max. Depth m
Transitional water category 1	< 5	< 1
Transitional water category 2	> 5	< 2
Transitional water category 3	< 30	> 2
Transitional water category 4	30 to 100	> 2
Transitional water category 5	> 100	> 2

Within a river or transitional water, categories according to the Tables 1 and 3 usually change from lower to higher ones according to longitudinal changes. But also within a relatively short stretch of a river or transitional water, categories can change from lower to higher ones and vice versa. In this case, the area itself of a river or transitional water, and not the whole river or transitional water, should be classified according to the given categories.

6 Fish sampling

6.1 General

Sampling methods are separated into methods by capture (6.2) and data sampling without fish capture (6.3). The principles are explained and the target water types or categories and fish species are given. The suitability of the methods is evaluated depending on the different water categories and their habitats as well as the target species. Target species are those species which can efficiently be caught by a particular method. Tables 4 to 6 provide a synopsis on sampling methods suitable or useful for the determination of the population parameters.

Methods not described below may be used in order to obtain additional information. They shall not be used exclusively.

6.2 Fish sampling by capture

6.2.1 Principle of fish capture

Fish may be caught by active gear (6.2.2 to 6.2.4) or passive (6.2.5 to 6.2.10). For all gear seasonal aspects shall be taken into account. The efficiency of active gear (electric fishing, seining, trawling) especially may vary considerably depending on the time (day/night) of use. Active gear is characterized by a comparatively short-term use, whereas passive ones normally are exposed over longer time periods. The fishing devices used shall be adapted to the local conditions as well as to the expected fish community. If it is intended to

Table 2 — Categories for lakes

Category	Presence or absence of a pelagic or profundal zone	Area km ²
Lake category 1	With or without a pelagic or profundal zone	< 0,5
Lake category 2	Without a pelagic or profundal zone	> 0,5
Lake category 3	With a pelagic and profundal zone	> 0,5

Table 3 — Categories for transitional waters

Category	Width m	max. Depth m
Transitional water category 1	< 5	< 1
Transitional water category 2	> 5	< 2
Transitional water category 3	< 30	> 2
Transitional water category 4	30 to 100	> 2
Transitional water category 5	> 100	> 2

Within a river or transitional water, categories according to the Tables 1 and 3 usually change from lower to higher ones according to longitudinal changes. But also within a relatively short stretch of a river or transitional water, categories can change from lower to higher ones and vice versa. In this case, the area itself of a river or transitional water, and not the whole river or transitional water, should be classified according to the given categories.

6 Fish sampling

6.1 General

Sampling methods are separated into methods by capture (6.2) and data sampling without fish capture (6.3). The principles are explained and the target water types or categories and fish species are given. The suitability of the methods is evaluated depending on the different water categories and their habitats as well as the target species. Target species are those species which can efficiently be caught by a particular method. Tables 4 to 6 provide a synopsis on sampling methods suitable or useful for the determination of the population parameters.

Methods not described below may be used in order to obtain additional information. They shall not be used exclusively.

6.2 Fish sampling by capture

6.2.1 Principle of fish capture

Fish may be caught by active gear (6.2.2 to 6.2.4) or passive (6.2.5 to 6.2.10). For all gear seasonal aspects shall be taken into account. The efficiency of active gear (electric fishing, seining, trawling) especially may vary considerably depending on the time (day/night) of use. Active gear is characterized by a comparatively short-term use, whereas passive ones normally are exposed over longer time periods. The fishing devices used shall be adapted to the local conditions as well as to the expected fish community. If it is intended to

Table 2 — Categories for lakes

Category	Presence or absence of a pelagic or profundal zone	Area km ²
Lake category 1	With or without a pelagic or profundal zone	< 0,5
Lake category 2	Without a pelagic or profundal zone	> 0,5
Lake category 3	With a pelagic and profundal zone	> 0,5

Table 3 — Categories for transitional waters

Category	Width m	max. Depth m
Transitional water category 1	< 5	< 1
Transitional water category 2	> 5	< 2
Transitional water category 3	< 30	> 2
Transitional water category 4	30 to 100	> 2
Transitional water category 5	> 100	> 2

Within a river or transitional water, categories according to the Tables 1 and 3 usually change from lower to higher ones according to longitudinal changes. But also within a relatively short stretch of a river or transitional water, categories can change from lower to higher ones and vice versa. In this case, the area itself of a river or transitional water, and not the whole river or transitional water, should be classified according to the given categories.

6 Fish sampling

6.1 General

Sampling methods are separated into methods by capture (6.2) and data sampling without fish capture (6.3). The principles are explained and the target water types or categories and fish species are given. The suitability of the methods is evaluated depending on the different water categories and their habitats as well as the target species. Target species are those species which can efficiently be caught by a particular method. Tables 4 to 6 provide a synopsis on sampling methods suitable or useful for the determination of the population parameters.

Methods not described below may be used in order to obtain additional information. They shall not be used exclusively.

6.2 Fish sampling by capture

6.2.1 Principle of fish capture

Fish may be caught by active gear (6.2.2 to 6.2.4) or passive (6.2.5 to 6.2.10). For all gear seasonal aspects shall be taken into account. The efficiency of active gear (electric fishing, seining, trawling) especially may vary considerably depending on the time (day/night) of use. Active gear is characterized by a comparatively short-term use, whereas passive ones normally are exposed over longer time periods. The fishing devices used shall be adapted to the local conditions as well as to the expected fish community. If it is intended to

Table 2 — Categories for lakes

Category	Presence or absence of a pelagic or profundal zone	Area km ²
Lake category 1	With or without a pelagic or profundal zone	< 0,5
Lake category 2	Without a pelagic or profundal zone	> 0,5
Lake category 3	With a pelagic and profundal zone	> 0,5

Table 3 — Categories for transitional waters

Category	Width m	max. Depth m
Transitional water category 1	< 5	< 1
Transitional water category 2	> 5	< 2
Transitional water category 3	< 30	> 2
Transitional water category 4	30 to 100	> 2
Transitional water category 5	> 100	> 2

Within a river or transitional water, categories according to the Tables 1 and 3 usually change from lower to higher ones according to longitudinal changes. But also within a relatively short stretch of a river or transitional water, categories can change from lower to higher ones and vice versa. In this case, the area itself of a river or transitional water, and not the whole river or transitional water, should be classified according to the given categories.

6 Fish sampling

6.1 General

Sampling methods are separated into methods by capture (6.2) and data sampling without fish capture (6.3). The principles are explained and the target water types or categories and fish species are given. The suitability of the methods is evaluated depending on the different water categories and their habitats as well as the target species. Target species are those species which can efficiently be caught by a particular method. Tables 4 to 6 provide a synopsis on sampling methods suitable or useful for the determination of the population parameters.

Methods not described below may be used in order to obtain additional information. They shall not be used exclusively.

6.2 Fish sampling by capture

6.2.1 Principle of fish capture

Fish may be caught by active gear (6.2.2 to 6.2.4) or passive (6.2.5 to 6.2.10). For all gear seasonal aspects shall be taken into account. The efficiency of active gear (electric fishing, seining, trawling) especially may vary considerably depending on the time (day/night) of use. Active gear is characterized by a comparatively short-term use, whereas passive ones normally are exposed over longer time periods. The fishing devices used shall be adapted to the local conditions as well as to the expected fish community. If it is intended to

Table 2 — Categories for lakes

Category	Presence or absence of a pelagic or profundal zone	Area km ²
Lake category 1	With or without a pelagic or profundal zone	< 0,5
Lake category 2	Without a pelagic or profundal zone	> 0,5
Lake category 3	With a pelagic and profundal zone	> 0,5

Table 3 — Categories for transitional waters

Category	Width m	max. Depth m
Transitional water category 1	< 5	< 1
Transitional water category 2	> 5	< 2
Transitional water category 3	< 30	> 2
Transitional water category 4	30 to 100	> 2
Transitional water category 5	> 100	> 2

Within a river or transitional water, categories according to the Tables 1 and 3 usually change from lower to higher ones according to longitudinal changes. But also within a relatively short stretch of a river or transitional water, categories can change from lower to higher ones and vice versa. In this case, the area itself of a river or transitional water, and not the whole river or transitional water, should be classified according to the given categories.

6 Fish sampling

6.1 General

Sampling methods are separated into methods by capture (6.2) and data sampling without fish capture (6.3). The principles are explained and the target water types or categories and fish species are given. The suitability of the methods is evaluated depending on the different water categories and their habitats as well as the target species. Target species are those species which can efficiently be caught by a particular method. Tables 4 to 6 provide a synopsis on sampling methods suitable or useful for the determination of the population parameters.

Methods not described below may be used in order to obtain additional information. They shall not be used exclusively.

6.2 Fish sampling by capture

6.2.1 Principle of fish capture

Fish may be caught by active gear (6.2.2 to 6.2.4) or passive (6.2.5 to 6.2.10). For all gear seasonal aspects shall be taken into account. The efficiency of active gear (electric fishing, seining, trawling) especially may vary considerably depending on the time (day/night) of use. Active gear is characterized by a comparatively short-term use, whereas passive ones normally are exposed over longer time periods. The fishing devices used shall be adapted to the local conditions as well as to the expected fish community. If it is intended to

Table 2 — Categories for lakes

Category	Presence or absence of a pelagic or profundal zone	Area km ²
Lake category 1	With or without a pelagic or profundal zone	< 0,5
Lake category 2	Without a pelagic or profundal zone	> 0,5
Lake category 3	With a pelagic and profundal zone	> 0,5

Table 3 — Categories for transitional waters

Category	Width m	max. Depth m
Transitional water category 1	< 5	< 1
Transitional water category 2	> 5	< 2
Transitional water category 3	< 30	> 2
Transitional water category 4	30 to 100	> 2
Transitional water category 5	> 100	> 2

Within a river or transitional water, categories according to the Tables 1 and 3 usually change from lower to higher ones according to longitudinal changes. But also within a relatively short stretch of a river or transitional water, categories can change from lower to higher ones and vice versa. In this case, the area itself of a river or transitional water, and not the whole river or transitional water, should be classified according to the given categories.

6 Fish sampling

6.1 General

Sampling methods are separated into methods by capture (6.2) and data sampling without fish capture (6.3). The principles are explained and the target water types or categories and fish species are given. The suitability of the methods is evaluated depending on the different water categories and their habitats as well as the target species. Target species are those species which can efficiently be caught by a particular method. Tables 4 to 6 provide a synopsis on sampling methods suitable or useful for the determination of the population parameters.

Methods not described below may be used in order to obtain additional information. They shall not be used exclusively.

6.2 Fish sampling by capture

6.2.1 Principle of fish capture

Fish may be caught by active gear (6.2.2 to 6.2.4) or passive (6.2.5 to 6.2.10). For all gear seasonal aspects shall be taken into account. The efficiency of active gear (electric fishing, seining, trawling) especially may vary considerably depending on the time (day/night) of use. Active gear is characterized by a comparatively short-term use, whereas passive ones normally are exposed over longer time periods. The fishing devices used shall be adapted to the local conditions as well as to the expected fish community. If it is intended to

Table 2 — Categories for lakes

Category	Presence or absence of a pelagic or profundal zone	Area km ²
Lake category 1	With or without a pelagic or profundal zone	< 0,5
Lake category 2	Without a pelagic or profundal zone	> 0,5
Lake category 3	With a pelagic and profundal zone	> 0,5

Table 3 — Categories for transitional waters

Category	Width m	max. Depth m
Transitional water category 1	< 5	< 1
Transitional water category 2	> 5	< 2
Transitional water category 3	< 30	> 2
Transitional water category 4	30 to 100	> 2
Transitional water category 5	> 100	> 2

Within a river or transitional water, categories according to the Tables 1 and 3 usually change from lower to higher ones according to longitudinal changes. But also within a relatively short stretch of a river or transitional water, categories can change from lower to higher ones and vice versa. In this case, the area itself of a river or transitional water, and not the whole river or transitional water, should be classified according to the given categories.

6 Fish sampling

6.1 General

Sampling methods are separated into methods by capture (6.2) and data sampling without fish capture (6.3). The principles are explained and the target water types or categories and fish species are given. The suitability of the methods is evaluated depending on the different water categories and their habitats as well as the target species. Target species are those species which can efficiently be caught by a particular method. Tables 4 to 6 provide a synopsis on sampling methods suitable or useful for the determination of the population parameters.

Methods not described below may be used in order to obtain additional information. They shall not be used exclusively.

6.2 Fish sampling by capture

6.2.1 Principle of fish capture

Fish may be caught by active gear (6.2.2 to 6.2.4) or passive (6.2.5 to 6.2.10). For all gear seasonal aspects shall be taken into account. The efficiency of active gear (electric fishing, seining, trawling) especially may vary considerably depending on the time (day/night) of use. Active gear is characterized by a comparatively short-term use, whereas passive ones normally are exposed over longer time periods. The fishing devices used shall be adapted to the local conditions as well as to the expected fish community. If it is intended to

Table 2 — Categories for lakes

Category	Presence or absence of a pelagic or profundal zone	Area km ²
Lake category 1	With or without a pelagic or profundal zone	< 0,5
Lake category 2	Without a pelagic or profundal zone	> 0,5
Lake category 3	With a pelagic and profundal zone	> 0,5

Table 3 — Categories for transitional waters

Category	Width m	max. Depth m
Transitional water category 1	< 5	< 1
Transitional water category 2	> 5	< 2
Transitional water category 3	< 30	> 2
Transitional water category 4	30 to 100	> 2
Transitional water category 5	> 100	> 2

Within a river or transitional water, categories according to the Tables 1 and 3 usually change from lower to higher ones according to longitudinal changes. But also within a relatively short stretch of a river or transitional water, categories can change from lower to higher ones and vice versa. In this case, the area itself of a river or transitional water, and not the whole river or transitional water, should be classified according to the given categories.

6 Fish sampling

6.1 General

Sampling methods are separated into methods by capture (6.2) and data sampling without fish capture (6.3). The principles are explained and the target water types or categories and fish species are given. The suitability of the methods is evaluated depending on the different water categories and their habitats as well as the target species. Target species are those species which can efficiently be caught by a particular method. Tables 4 to 6 provide a synopsis on sampling methods suitable or useful for the determination of the population parameters.

Methods not described below may be used in order to obtain additional information. They shall not be used exclusively.

6.2 Fish sampling by capture

6.2.1 Principle of fish capture

Fish may be caught by active gear (6.2.2 to 6.2.4) or passive (6.2.5 to 6.2.10). For all gear seasonal aspects shall be taken into account. The efficiency of active gear (electric fishing, seining, trawling) especially may vary considerably depending on the time (day/night) of use. Active gear is characterized by a comparatively short-term use, whereas passive ones normally are exposed over longer time periods. The fishing devices used shall be adapted to the local conditions as well as to the expected fish community. If it is intended to

Table 2 — Categories for lakes

Category	Presence or absence of a pelagic or profundal zone	Area km ²
Lake category 1	With or without a pelagic or profundal zone	< 0,5
Lake category 2	Without a pelagic or profundal zone	> 0,5
Lake category 3	With a pelagic and profundal zone	> 0,5

Table 3 — Categories for transitional waters

Category	Width m	max. Depth m
Transitional water category 1	< 5	< 1
Transitional water category 2	> 5	< 2
Transitional water category 3	< 30	> 2
Transitional water category 4	30 to 100	> 2
Transitional water category 5	> 100	> 2

Within a river or transitional water, categories according to the Tables 1 and 3 usually change from lower to higher ones according to longitudinal changes. But also within a relatively short stretch of a river or transitional water, categories can change from lower to higher ones and vice versa. In this case, the area itself of a river or transitional water, and not the whole river or transitional water, should be classified according to the given categories.

6 Fish sampling

6.1 General

Sampling methods are separated into methods by capture (6.2) and data sampling without fish capture (6.3). The principles are explained and the target water types or categories and fish species are given. The suitability of the methods is evaluated depending on the different water categories and their habitats as well as the target species. Target species are those species which can efficiently be caught by a particular method. Tables 4 to 6 provide a synopsis on sampling methods suitable or useful for the determination of the population parameters.

Methods not described below may be used in order to obtain additional information. They shall not be used exclusively.

6.2 Fish sampling by capture

6.2.1 Principle of fish capture

Fish may be caught by active gear (6.2.2 to 6.2.4) or passive (6.2.5 to 6.2.10). For all gear seasonal aspects shall be taken into account. The efficiency of active gear (electric fishing, seining, trawling) especially may vary considerably depending on the time (day/night) of use. Active gear is characterized by a comparatively short-term use, whereas passive ones normally are exposed over longer time periods. The fishing devices used shall be adapted to the local conditions as well as to the expected fish community. If it is intended to

Table 2 — Categories for lakes

Category	Presence or absence of a pelagic or profundal zone	Area km ²
Lake category 1	With or without a pelagic or profundal zone	< 0,5
Lake category 2	Without a pelagic or profundal zone	> 0,5
Lake category 3	With a pelagic and profundal zone	> 0,5

Table 3 — Categories for transitional waters

Category	Width m	max. Depth m
Transitional water category 1	< 5	< 1
Transitional water category 2	> 5	< 2
Transitional water category 3	< 30	> 2
Transitional water category 4	30 to 100	> 2
Transitional water category 5	> 100	> 2

Within a river or transitional water, categories according to the Tables 1 and 3 usually change from lower to higher ones according to longitudinal changes. But also within a relatively short stretch of a river or transitional water, categories can change from lower to higher ones and vice versa. In this case, the area itself of a river or transitional water, and not the whole river or transitional water, should be classified according to the given categories.

6 Fish sampling

6.1 General

Sampling methods are separated into methods by capture (6.2) and data sampling without fish capture (6.3). The principles are explained and the target water types or categories and fish species are given. The suitability of the methods is evaluated depending on the different water categories and their habitats as well as the target species. Target species are those species which can efficiently be caught by a particular method. Tables 4 to 6 provide a synopsis on sampling methods suitable or useful for the determination of the population parameters.

Methods not described below may be used in order to obtain additional information. They shall not be used exclusively.

6.2 Fish sampling by capture

6.2.1 Principle of fish capture

Fish may be caught by active gear (6.2.2 to 6.2.4) or passive (6.2.5 to 6.2.10). For all gear seasonal aspects shall be taken into account. The efficiency of active gear (electric fishing, seining, trawling) especially may vary considerably depending on the time (day/night) of use. Active gear is characterized by a comparatively short-term use, whereas passive ones normally are exposed over longer time periods. The fishing devices used shall be adapted to the local conditions as well as to the expected fish community. If it is intended to

Table 2 — Categories for lakes

Category	Presence or absence of a pelagic or profundal zone	Area km ²
Lake category 1	With or without a pelagic or profundal zone	< 0,5
Lake category 2	Without a pelagic or profundal zone	> 0,5
Lake category 3	With a pelagic and profundal zone	> 0,5

Table 3 — Categories for transitional waters

Category	Width m	max. Depth m
Transitional water category 1	< 5	< 1
Transitional water category 2	> 5	< 2
Transitional water category 3	< 30	> 2
Transitional water category 4	30 to 100	> 2
Transitional water category 5	> 100	> 2

Within a river or transitional water, categories according to the Tables 1 and 3 usually change from lower to higher ones according to longitudinal changes. But also within a relatively short stretch of a river or transitional water, categories can change from lower to higher ones and vice versa. In this case, the area itself of a river or transitional water, and not the whole river or transitional water, should be classified according to the given categories.

6 Fish sampling

6.1 General

Sampling methods are separated into methods by capture (6.2) and data sampling without fish capture (6.3). The principles are explained and the target water types or categories and fish species are given. The suitability of the methods is evaluated depending on the different water categories and their habitats as well as the target species. Target species are those species which can efficiently be caught by a particular method. Tables 4 to 6 provide a synopsis on sampling methods suitable or useful for the determination of the population parameters.

Methods not described below may be used in order to obtain additional information. They shall not be used exclusively.

6.2 Fish sampling by capture

6.2.1 Principle of fish capture

Fish may be caught by active gear (6.2.2 to 6.2.4) or passive (6.2.5 to 6.2.10). For all gear seasonal aspects shall be taken into account. The efficiency of active gear (electric fishing, seining, trawling) especially may vary considerably depending on the time (day/night) of use. Active gear is characterized by a comparatively short-term use, whereas passive ones normally are exposed over longer time periods. The fishing devices used shall be adapted to the local conditions as well as to the expected fish community. If it is intended to

Table 2 — Categories for lakes

Category	Presence or absence of a pelagic or profundal zone	Area km ²
Lake category 1	With or without a pelagic or profundal zone	< 0,5
Lake category 2	Without a pelagic or profundal zone	> 0,5
Lake category 3	With a pelagic and profundal zone	> 0,5

Table 3 — Categories for transitional waters

Category	Width m	max. Depth m
Transitional water category 1	< 5	< 1
Transitional water category 2	> 5	< 2
Transitional water category 3	< 30	> 2
Transitional water category 4	30 to 100	> 2
Transitional water category 5	> 100	> 2

Within a river or transitional water, categories according to the Tables 1 and 3 usually change from lower to higher ones according to longitudinal changes. But also within a relatively short stretch of a river or transitional water, categories can change from lower to higher ones and vice versa. In this case, the area itself of a river or transitional water, and not the whole river or transitional water, should be classified according to the given categories.

6 Fish sampling

6.1 General

Sampling methods are separated into methods by capture (6.2) and data sampling without fish capture (6.3). The principles are explained and the target water types or categories and fish species are given. The suitability of the methods is evaluated depending on the different water categories and their habitats as well as the target species. Target species are those species which can efficiently be caught by a particular method. Tables 4 to 6 provide a synopsis on sampling methods suitable or useful for the determination of the population parameters.

Methods not described below may be used in order to obtain additional information. They shall not be used exclusively.

6.2 Fish sampling by capture

6.2.1 Principle of fish capture

Fish may be caught by active gear (6.2.2 to 6.2.4) or passive (6.2.5 to 6.2.10). For all gear seasonal aspects shall be taken into account. The efficiency of active gear (electric fishing, seining, trawling) especially may vary considerably depending on the time (day/night) of use. Active gear is characterized by a comparatively short-term use, whereas passive ones normally are exposed over longer time periods. The fishing devices used shall be adapted to the local conditions as well as to the expected fish community. If it is intended to

Table 2 — Categories for lakes

Category	Presence or absence of a pelagic or profundal zone	Area km ²
Lake category 1	With or without a pelagic or profundal zone	< 0,5
Lake category 2	Without a pelagic or profundal zone	> 0,5
Lake category 3	With a pelagic and profundal zone	> 0,5

Table 3 — Categories for transitional waters

Category	Width m	max. Depth m
Transitional water category 1	< 5	< 1
Transitional water category 2	> 5	< 2
Transitional water category 3	< 30	> 2
Transitional water category 4	30 to 100	> 2
Transitional water category 5	> 100	> 2

Within a river or transitional water, categories according to the Tables 1 and 3 usually change from lower to higher ones according to longitudinal changes. But also within a relatively short stretch of a river or transitional water, categories can change from lower to higher ones and vice versa. In this case, the area itself of a river or transitional water, and not the whole river or transitional water, should be classified according to the given categories.

6 Fish sampling

6.1 General

Sampling methods are separated into methods by capture (6.2) and data sampling without fish capture (6.3). The principles are explained and the target water types or categories and fish species are given. The suitability of the methods is evaluated depending on the different water categories and their habitats as well as the target species. Target species are those species which can efficiently be caught by a particular method. Tables 4 to 6 provide a synopsis on sampling methods suitable or useful for the determination of the population parameters.

Methods not described below may be used in order to obtain additional information. They shall not be used exclusively.

6.2 Fish sampling by capture

6.2.1 Principle of fish capture

Fish may be caught by active gear (6.2.2 to 6.2.4) or passive (6.2.5 to 6.2.10). For all gear seasonal aspects shall be taken into account. The efficiency of active gear (electric fishing, seining, trawling) especially may vary considerably depending on the time (day/night) of use. Active gear is characterized by a comparatively short-term use, whereas passive ones normally are exposed over longer time periods. The fishing devices used shall be adapted to the local conditions as well as to the expected fish community. If it is intended to

Table 2 — Categories for lakes

Category	Presence or absence of a pelagic or profundal zone	Area km ²
Lake category 1	With or without a pelagic or profundal zone	< 0,5
Lake category 2	Without a pelagic or profundal zone	> 0,5
Lake category 3	With a pelagic and profundal zone	> 0,5

Table 3 — Categories for transitional waters

Category	Width m	max. Depth m
Transitional water category 1	< 5	< 1
Transitional water category 2	> 5	< 2
Transitional water category 3	< 30	> 2
Transitional water category 4	30 to 100	> 2
Transitional water category 5	> 100	> 2

Within a river or transitional water, categories according to the Tables 1 and 3 usually change from lower to higher ones according to longitudinal changes. But also within a relatively short stretch of a river or transitional water, categories can change from lower to higher ones and vice versa. In this case, the area itself of a river or transitional water, and not the whole river or transitional water, should be classified according to the given categories.

6 Fish sampling

6.1 General

Sampling methods are separated into methods by capture (6.2) and data sampling without fish capture (6.3). The principles are explained and the target water types or categories and fish species are given. The suitability of the methods is evaluated depending on the different water categories and their habitats as well as the target species. Target species are those species which can efficiently be caught by a particular method. Tables 4 to 6 provide a synopsis on sampling methods suitable or useful for the determination of the population parameters.

Methods not described below may be used in order to obtain additional information. They shall not be used exclusively.

6.2 Fish sampling by capture

6.2.1 Principle of fish capture

Fish may be caught by active gear (6.2.2 to 6.2.4) or passive (6.2.5 to 6.2.10). For all gear seasonal aspects shall be taken into account. The efficiency of active gear (electric fishing, seining, trawling) especially may vary considerably depending on the time (day/night) of use. Active gear is characterized by a comparatively short-term use, whereas passive ones normally are exposed over longer time periods. The fishing devices used shall be adapted to the local conditions as well as to the expected fish community. If it is intended to